

## Multiple Sclerosis Enrollment Form Phone: 1-877-437-9012 Fax 1-877-309-0687

□ New to Therapy □ Current Therapy

Patient Information	Clinical Information and Prescription
Patient Name:	Diagnosis:  □G35 Multiple Sclerosis  □Other:    Type:  □RRMS  □PPMS  □SPMS  □Clinically definite MS    Date of Diagnosis or Years with Disease:   Patients Allergies:    Patient Weight: kgs  or lbs  (please indicate) Patient Height: inches  or cm.  (please indicate)    Prior Treatments with duration and reason for discontinuation
Home Phone:  Work Phone:    Cell Phone:  E-mail:    Please attach copy of front and back of patient's prescription ins. card(s) if applicable    Insurance Company Name:    Insurance Company Phone:	□ Aubagio® (teriflunomide)  □ 7 mg  □ 14 mg  One tablet by mouth once daily    □ Avonex® (interferon -β1a)  Please indicate dosage form:  □ Pen  □ Prefilled Syringe  □ Single Dose Vials    □ Initial titration:  Inject IM as follows:  7.5mcg on week 1; 15mcg IM on week 2; 22.5mcg IM on week 3; then 30mcg IM on week 4 and once weekly thereafter.  Dispense PFS with AVOSTARTGRIP™ Titration device.    □ Inject 30mcg IM once a week  □
Policyholder: Policy holder Employer: Relationship to Patient:	□ Betaseron® (interferon-β1b)  □ Initial titration: 62.5mcg increasing to 250 mcg SC every other day over 6 weeks as directed    □ Inject 250mcg SC every other day    □ Dalfampridine (generic for Ampyra®) 10 mg tablet by mouth every 12 hours    □ Extavia® (interferon-β1b)  □ Initial titration: 62.5mcg increasing to 250 mcg SC every other day over 6 weeks as directed    □ Inject 250mcg SC every other day  □ Initial titration: 62.5mcg increasing to 250 mcg SC every other day over 6 weeks as directed
Prescriber Information	□ Inject 250mcg SC every other day □ Gilenya® (fingolimod) □ 0.5mg □ 0.25 mg Take 1 capsule by mouth once daily Date of First Dose Observation://
Practice/ Organization Name: Physician Name: Address: City:State:Zip: Phone#:Fax#: DEA#NPI# License#:Nedicaid UPIN#: Physician Specialty: Physician Specialty: Date Shipment Needed: _//Attn: Ship to:PatientPrescriberInfusion Clinic Shipment Address: City:State:Zip: If shipped to the physician's office, physician accepts on behalf of patient for administration in office.	□ Glatiramer acetate OR □ Copaxone® OR □ Glatopa® □ Inject 40mg SC three times weekly OR □ Inject 20mg SC daily    □ Mayzent® (siponimod) *Please note: Elixir Specialty proudly services Mayzent patients but Novartis AlongsideMS    requires all initial referrals to route through www.mayzent.com or 877-MAYZENT (629-9368)    □ Novantrone® (mitoxantrone) 12 mg/m2 IV every 3 months.    □ Ocrevus® (ocrelizumab) □ Initial titration: 300 mg IV infusion as a single dose, followed by a 2 <sup>nd</sup> 300 mg IV infusion 2 weeks later    □ Subsequent infusions: 600 mg IV infusion every 6 months (1 <sup>st</sup> dose due 6 months after infusion 1 of the initial dose)    □ Plegridy® (pegylated interferon-β1b) Please indicate dosage form: □ Pen □ Prefilled Syringe    □ Starter Kit Needed: Inject 63mcg SC on day 1, 94mcg on day 15, then 125mcg every 14 days thereafter    □ Inject 125mcg SC every 14 days    □ Rebif® (interferon-β1a) Please indicate dosage form: □ Prefilled Syringe □ Rebidose    □ Titration to 22mcg: Inject SC as follows: 4.4mcg 3X/wk on weeks 1 & 2; 11mcg 3X/wk on weeks 3 & 4; then 22mcg 3x/wk    □ Titration to 44mcg: Inject SC as follows: 8.8mcg 3X/wk on weeks 1 & 2; 22mcg 3X/wk on weeks 3 & 4; then 44mcg 3x/wk    □ Tecfidera® (dimethyl fumarate) □ Starter Pack: Take 120mg orally twice a day for 7 days, then 240mg twice a day thereafter    □ Take 1 capsule (240mg) orally twice daily □ Other:
Privacy & Confidentiality of Information Notice: This communication may contain non- public, confidential, or legally privileged information intended for the sole use of the designated recipient(s). If you are not the intended recipient, or have received this communication in error, please notify the sender immediately by reply email or by telephone at the number stated above and delete all copies of this communication, including any attachments, without reading them or saving them to disk. If you are the intended recipient, you must secure the contents of this communication in accordance with all applicable state or federal requirements related to the privacy and confidentiality of information, including the HIPAA Privacy guidelines.	□  Vumerity® (diroximel fumarate)  □  Starter Pack: Take 231mg orally twice a day for 7 days, then 462mg twice a day thereafter    □  Take 2 capsules (462mg) orally twice daily  □  Other: