

Patient Information	Prescriber Information
Patient Name: _____ Date of Birth: _____ Gender: _____ SS# _____ Address: _____ City: _____ State: _____ Zip: _____ Home Phone: _____ Work Phone: _____ Cell Phone: _____ E-mail: _____ Patient Weight: _____ kg /lbs Patient Height: _____ in /cm Please attach copy of front and back of patient's prescription ins. card(s) if applicable Insurance Company Name: _____ Insurance Company Phone: _____ Policy holder: _____ Policy holder Employer: _____ Relationship to Patient: _____ ID# _____ Group# _____ RxBIN: _____ RxPCN: _____	Practice/ Organization Name: _____ Physician Name: _____ Contact Person: _____ Contact's Phone: _____ Address: _____ City: _____ State: _____ Zip: _____ Office Phone#: _____ Office Fax#: _____ DEA# _____ NPI# _____ License#: _____ Medicaid UPIN#: _____ Physician Specialty: _____ Date Shipment Needed: _____ Ship to: <input type="checkbox"/> Patient <input type="checkbox"/> Prescriber <input type="checkbox"/> Other _____ Shipment Address: _____ Attn: _____ City: _____ State: _____ Zip: _____ <i>If shipped to the physician's office, physician accepts on behalf of patient for administration in office.</i>

Diagnosis and Clinical Information

Diagnosis (ICD-10): J82 Pulmonary eosinophilia J45.4 Moderate Persistent Asthma J45.5 Severe Persistent Asthma D72.119 Hypereosinophilic syndrome (HES)
 M30.1 Eosinophilic Granulomatosis with Polyangiitis (EGPA) J33.0 Polyp of the nasal cavity J33.1 Polypoid sinus degeneration J33.8 Other polyp of sinus
 J33.9 Nasal polyp, unspecified K20.0 Eosinophilic esophagitis (EOE) Other: _____ Description: _____

Date of Diagnosis or Years with Disease: _____

Patients Allergies: _____ Latex allergy: Yes No

Comorbidities: _____

Lab Results (attach if available) : Past positive skin or RAST test to perennial aeroallergen Pre-treatment: Serum IgE level _____ IU/ML Testing date _____
 Serum eosinophils _____ cells/mcL; Sputum eosinophils _____ Testing date _____

Previous and Current Medication Use:

Short-acting beta agonist Long-acting beta agonist Antihistamines Decongestants Immunotherapy Inhaled corticosteroids Leukotriene modifiers
 Oral steroids Nasal steroids Other: _____

Expected First Dose Date: _____ Injection training needed: Yes No EpiPen on hand for Patient: Yes No

Shipment to Patient Yes No Shipment to Provider: Yes No Shipment Address: _____

Privacy & Confidentiality of Information Notice: This communication may contain non-public, confidential, or legally privileged information intended for the sole use of the designated recipient(s). If you are not the intended recipient, or have received this communication in error, please notify the sender immediately by reply email or by telephone at the number stated above and delete all copies of this communication, including any attachments, without reading them or saving them to disk. If you are the intended recipient, you must secure the contents of this communication in accordance with all applicable state or federal requirements related to the privacy and confidentiality of information, including the HIPAA Privacy guidelines.

<input type="checkbox"/> New to Therapy
<input type="checkbox"/> Current Therapy

Patient Name: _____ **Patient Date of Birth:** _____

Prescriber Name: _____ **Phone:** _____ **Date:** _____

Prescriber Address: _____

Dupixent® (dupilumab) 100 mg pre-filled syringe 200 mg pre-filled syringe 200 mg pen-injector 300 mg pre-filled syringe 300 mg pen-injector Patient weight: _____

Starter: Inject 2 syringes subcutaneously on day 1 then 1 syringe on day 15

Maintenance: Injection 1 syringe subcutaneously every 2 weeks

Nucala® (mepolizumab) 100 mg vial 100 mg pre-filled syringe 100 mg auto-injector **Cinqair®** (reslizumab, 100 mg/10mL single-use vial)

Inject 100 mg subcutaneously once every 4 weeks. 3 mg/kg Other: _____

To be given by IV infusion once every 4 weeks by prescriber as directed.

Xolair® (omalizumab) 75mg pre-filled syringe 150mg pre-filled syringe 150mg vial

To be injected subcutaneously once every 4 weeks as directed:

75 mg 150 mg 225 mg 300 mg Other: _____

To be injected subcutaneously once every 2 weeks as directed:

225 mg 300 mg 375 mg Other: _____

Sterile Water for Injection

To be used as directed to reconstitute **Xolair®** **Nucala®** vials

Quantity: _____ vial(s) Refills: _____

EpiPen **EpiPen Jr**

Use as directed for anaphylaxis

Quantity: 2 Pens Refills: _____

Quantity Prescribed: QS 30 days Other: _____

Refills Authorized: 0 1 2 3 6 11 Other: _____

Dispense: 28 day supply of medication with 0 1 2 Other: _____ Refills

If Brand required "Dispense as Written" must be handwritten

Prescriber Signature: X _____ **Date:** _____

Privacy & Confidentiality of Information Notice: This communication may contain non-public, confidential, or legally privileged information intended for the sole use of the designated recipient(s). If you are not the intended recipient, or have received this communication in error, please notify the sender immediately by reply email or by telephone at the number stated above and delete all copies of this communication, including any attachments, without reading them or saving them to disk. If you are the intended recipient, you must secure the contents of this communication in accordance with all applicable state or federal requirements related to the privacy and confidentiality of information, including the HIPAA Privacy guidelines.